

ALCOHOL KILLS... Bacteria

Reducing Ineffective and Counterproductive Use of Hand Sanitizers

Members: Luke Goodman (RBE), Brittany Jones (BBT), David Karbassi (BBT), Angela Wood (BBT/PW) Advisors: Jill Rulfs (BBT), Helen Vassallo (MGMT)

Need

To promote better hand hygiene in the WPI community, thereby decreasing the spread of disease.

Approach

- Design a survey to gather information regarding use and understanding of hand sanitizer
- Distribute the survey to the WPI student body through email
- > Distribute "info cards" to raise awareness

Literature Review

65% 60%

Alcohol Concentration by Brand*



*Alcohol Content Specified on Product

- > Alcohol-based sanitizers must contain at least 60% alcohol to be effective
- > QACs such as Benzethonium and Benzalkonium Chloride are antimicrobial agents found in hand sanitizers, surface cleaners, etc.
- Research in progress shows bacteria developing resistance to QACs and other agents

Data

Chart I: Frequency of Use (Times per Day)




Chart II: Awareness of Different

Active Ingredients

59.1%/

Aware


■ Not Aware

0.9%

40.0%

70%

Chart III: Preferred Active Ingredients




Alcohol

■ Non-alcohol

■ No preference

Chart IV: Effectiveness of Different Alcohol Concentrations



Outcome

- > Total responses: 467
- > Important Findings:
 - Chart I: The WPI community uses hand sanitizer, but not excessively
 - Charts II and III: Many are unaware of or simply don't consider the active ingredients in the sanitizer they use
 - Chart IV: Similarly, many are unaware that a minimum concentration of alcohol (60%) is required to be effective

Considerations

- > Hawthorne Effect: questions designed to avoid this, but it may have influenced the results regardless
- The "Preferred Active Ingredients" chart excludes the "No Response" category




Image I



Image II

Image Sources:

I: http://content9.eol.org/content/2009/11/25/03/07773_large.jpg II: http://farm3.static.flickr.com/2739/4402085217_31183b8c74.jpg

Acknowledgements

We would like to thank Jill Rulfs, Helen Vassallo, and the ATC for their help.

References

Aiello, Allison E. et al. "Effect of Hand Hygiene on Infectious Disease Risk in the Community Setting: A Meta-Analysis." Framing Americans Use Hand Sanitizers for Peace of Mind but Don't Know Leading Hand Sanitizers Stop Working After Two Minutes.' PR Newswire 4 Nov. 2010. Academic OneFile. Web. 7 Dec. 2010

Am J Infect Control. 2007 May; 35(4): 231-236.

Braoudaki, M., and A. C. Hilton. "Adaptive Resistance to Biocides in Salmonella enterica and Escherichia coli O157 and Cross-Resistance to Antimicrobial Agents." Journal of Clinical Microbiology. (2004): 73-78. Print

Containing Quaternary Ammonium Compounds and Drug Carson, Robyn T. "Use of Antibacterial Consumer Products Resistance in the Community." Journal of Antimicrobial Chemotherapy. (2008): 1160-61. Print.

'Health Buzz: FDA Will Review Academic Safety of Antibacterial in Soap." Wall Street Journal 12 April 2010. LexisNexis

"The bugs stop here." Men's Health Mar. 2010: 64. Academic OneFile. Web. 7 Dec. 2010.

Background from: http://www.public-domain-image.com/backgrounds/slides/abstract-waves-on-a-blue-background.html